

REAL ALGEBRAIC GEOMETRY LECTURE NOTES
(25: 26/01/10)

SALMA KUHLMANN

CONTENTS

1.	Introduction	1
2.	Pseudo-convergence and maximality	1
3.	Pseudo-limits	3
4.	Cofinal subsets	4

1. INTRODUCTION

Our aim for this and next lecture is to complete the proof of Hahn's embedding Theorem:

Let (V, v) be a Q -valued vector space with $S(V) = [\Gamma, B(\gamma)]$.
 Let $\{x_i : i \in I\} \subset V$ be maximal valuation independent and

$$h: V_0 = (\langle \{x_i : i \in I\} \rangle, v) \xrightarrow{\sim} \left(\bigsqcup_{\gamma \in \Gamma} B(\gamma), v_{\min} \right).$$

Then h extends to a valuation preserving embedding (i.e. an isomorphism onto a valued subspace)

$$\tilde{h}: (V, v) \hookrightarrow (\mathbb{H}_{\gamma \in \Gamma} B(\gamma), v_{\min}).$$

The picture is the following:

$$\begin{array}{ccc} (V, v) & \xhookrightarrow{\tilde{h}} & (\mathbb{H}_{\gamma \in \Gamma} B(\gamma), v_{\min}) \\ \text{immediate} \Big| & & \Big| \text{immediate} \\ (V_0, v) & \xrightarrow[\sim]{h} & \left(\bigsqcup_{\gamma \in \Gamma} B(\gamma), v_{\min} \right) \end{array}$$

2. PSEUDO-CONVERGENCE AND MAXIMALITY

Definition 2.1. A valued Q -vector space (V, v) is said to be **maximally valued** if it admits no proper immediate extension.

Definition 2.2. A well ordered set $S = \{a_\rho : \rho \in \lambda\} \subset V$ without a last element is said to be **pseudo-convergent** (or **pseudo-Cauchy**) if for every $\rho < \sigma < \tau$ we have

$$v(a_\sigma - a_\rho) < v(a_\tau - a_\sigma).$$

Example 2.3.

(a) Let $V = (H_{\mathbb{N}_0} \mathbb{R}, v_{\min})$, where $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. An element $s \in V$ can be viewed as a function $s: \mathbb{N}_0 \rightarrow \mathbb{R}$. Consider

$$a_0 = (1, 0, 0, 0, 0 \dots)$$

$$a_1 = (1, 1, 0, 0, 0 \dots)$$

$$a_2 = (1, 1, 1, 0, 0 \dots)$$

\vdots

The sequence $\{a_n : n \in \mathbb{N}_0\} \subset V$ is pseudo-Cauchy.

(b) Take V as above and $s \in V$ with

$$\text{support}(s) = \mathbb{N}_0,$$

i.e. $s_i := s(i) \neq 0 \forall i \in \mathbb{N}_0$. Define the sequence

$$b_0 = (s_0, 0, 0, 0, 0 \dots)$$

$$b_1 = (s_0, s_1, 0, 0, 0 \dots)$$

$$b_2 = (s_0, s_1, s_2, 0, 0 \dots)$$

\vdots

For every $l < m < n \in \mathbb{N}_0$, we have

$$l + 1 = v_{\min}(b_m - b_l) < v_{\min}(b_n - b_m) = m + 1.$$

Therefore $\{b_n : n \in \mathbb{N}_0\} \subset V$ is pseudo-Cauchy.

Lemma 2.4. If $S = \{a_\rho\}_{\rho \in \lambda}$ is pseudo-convergent then

(i) either $v(a_\rho) < v(a_\sigma)$ for all $\rho < \sigma \in \lambda$,

(ii) or $\exists \rho_0 \in \lambda$ such that $v(a_\rho) = v(a_\sigma) \forall \rho, \sigma \geq \rho_0$.

Proof. Assume (i) does not hold, i.e. $v(a_\rho) \geq v(a_\sigma)$ for some $\rho < \sigma$. Then we claim that

$$v(a_\tau) = v(a_\sigma) \quad \forall \tau > \sigma.$$

Otherwise, $v(a_\tau - a_\sigma) = \min\{v(a_\tau), v(a_\sigma)\} \leq v(a_\sigma)$.

But $v(a_\sigma - a_\rho) \geq v(a_\sigma)$, contradicting 2.2. \square

Notation 2.5. In case (ii) define

$$\text{Ult } S := v(a_{\rho_0}) = v(a_\rho) \quad \forall \rho \geq \rho_0.$$

Lemma 2.6. *If $\{a_\rho\}$ is pseudo-convergent then for all $\rho < \sigma$ we have*

$$v(a_\sigma - a_\rho) = v(a_{\rho+1} - a_\rho).$$

Proof. We may assume $\sigma > \rho + 1$ (so $\rho < \rho + 1 < \sigma$). From

$$v(a_{\rho+1} - a_\rho) < v(a_\sigma - a_{\rho+1})$$

and the identity

$$a_\sigma - a_\rho = (a_\sigma - a_{\rho+1}) + (a_{\rho+1} - a_\rho),$$

we deduce that

$$\begin{aligned} v(a_\sigma - a_\rho) &= \min(v(a_\sigma - a_{\rho+1}), v(a_{\rho+1} - a_\rho)) \\ &= v(a_{\rho+1} - a_\rho). \end{aligned}$$

□

Notation 2.7.

$$\begin{aligned} \gamma_\rho &:= v(a_{\rho+1} - a_\rho) \\ &= v(a_\sigma - a_\rho) \quad \forall \sigma > \rho. \end{aligned}$$

Remark 2.8. Since $\rho < \rho + 1 < \rho + 2$, we have $\gamma_\rho < \gamma_{\rho+1}$ for all ρ .

3. PSEUDO-LIMITS

Definition 3.1. Let $S = \{a_\rho\}$ be a pseudo-convergent set. We say that $x \in V$ is a **pseudo-limit** of S if

$$v(x - a_\rho) = \gamma_\rho \quad \text{for all } \rho.$$

Remark 3.2.

(i) If $v(a_\rho) < v(a_\sigma)$ for $\rho < \sigma$, then $x = 0$ is a pseudo-limit.

(ii) If 0 is not a pseudo-limit and x is a pseudo-limit, then $v(x) = \text{Ult } S$.

Example 3.3.

(a) In Example 2.3(a), the constant function 1:

$$a = (1, 1, \dots)$$

is a pseudo-limit of the sequence $\{a_n\}_{n \in \mathbb{N}_0}$.

(b) In Example 2.3(b), s is a pseudo-limit of $\{b_n\}_{n \in \mathbb{N}_0}$.

Definition 3.4. (V, v) is **pseudo-complete** if every pseudo-convergent sequence has a pseudo-limit in V .

Definition 3.5. Let $S = \{a_\rho\}$ be pseudo-convergent. The **breadth** (*Breite*) B of S is defined to be the following subset of V :

$$B(S) = \{y \in V : v(y) > \gamma_\rho \forall \rho\}.$$

Lemma 3.6. Let $\{a_\rho\}$ be pseudo-convergent with breadth B and let $x \in V$ be a pseudo-limit. Then an element of V is a pseudo-limit of $\{a_\rho\}$ if and only if it is of the form $x + y$ with $y \in B$.

Proof.

(\Rightarrow) Let z be another pseudo-limit of $\{a_\rho\}$. It follows from

$$x - z = (x - a_\rho) - (z - a_\rho)$$

that

$$v(x - z) \geq \min\{v(x - a_\rho), v(z - a_\rho)\} = \gamma_\rho \quad \forall \rho.$$

Since γ_ρ is increasing, it follows that $v(x - z) > \gamma_\rho$ for all ρ . So $z \in B$ as required.

(\Leftarrow) If $y \in B$ then $v(y) > \gamma_\rho = v(x - a_\rho)$ for all ρ . Then

$$v(x + y - a_\rho) = v(x - a_\rho + y) = \min\{v(x - a_\rho), v(y)\} = \gamma_\rho \quad \forall \rho.$$

□

4. COFINAL SUBSETS

Definition 4.1. Let Γ be a totally ordered set. A subset $A \subset \Gamma$ is cofinal in Γ if

$$\forall \gamma \in \Gamma \exists a \in A \text{ with } \gamma \leq a.$$

Example 4.2. If $\Gamma = [0, 1] \subset \mathbb{R}$, then for instance $A = \{1\}$ is cofinal in Γ .

Lemma 4.3. Let Γ be a totally ordered set. Then there is a well ordered cofinal subset $A \subset \Gamma$. Moreover if Γ has no last element, then also A has no last element.