

**REAL ALGEBRAIC GEOMETRY LECTURE NOTES**  
**(08: 12/11/09)**

SALMA KUHLMANN

CONTENTS

1. Real closure	1
2. Order preserving extensions	2

1. REAL CLOSURE

**Definition 1.1.** Let  $(K, P)$  be an ordered field.  $R$  is a real closure of  $(K, P)$  if

- (1)  $R$  is real closed,
- (2)  $R \supseteq K$ ,  $R|K$  is algebraic,
- (3)  $P = \sum R^2 \cap K$  (i.e. the order on  $K$  is the restriction of the unique order  $R$  to  $K$ ).

**Theorem 1.2.** *Every ordered field  $(K, P)$  has a real closure.*

*Proof.* Apply Zorn's Lemma to

$$\mathcal{L} := \{(L, Q) : L|K \text{ algebraic, } Q \cap K = P\}.$$

□

**Proposition 1.3.** *(Corollary to Sturm's Theorem) Let  $K$  be a field. Let  $R_1, R_2$  be two real closed fields such that*

$$K \subseteq R_1 \quad \text{and} \quad K \subseteq R_2$$

*with*

$$P := K \cap \sum R_1^2 = K \cap \sum R_2^2$$

*(i.e.  $R_1$  and  $R_2$  induce the same ordering  $P$  on  $K$ ).*

*Let  $f(x) \in K[x]$ ; then the number of roots of  $f(x)$  in  $R_1$  is equal to the number of roots of  $f(x)$  in  $R_2$ .*

## 2. ORDER PRESERVING EXTENSIONS

**Proposition 2.1.** *Let  $(K, P)$  be an ordered field. Let  $R$  be a real closed field containing  $(K, P)$ . Let  $K \subseteq L \subseteq R$  be such that  $[L : K] < \infty$ . Let  $S$  be a real closed field with*

$$\varphi: (K, P) \hookrightarrow (S, \sum S^2)$$

*an order preserving embedding. Then  $\varphi$  extends to an order preserving embedding*

$$\psi: (L, \sum R^2 \cap L) \hookrightarrow (S, \sum S^2).$$

*Proof.* We recall that if  $(K, P)$  and  $(L, Q)$  are ordered fields, a field homomorphism  $\varphi: K \rightarrow L$  is called **order preserving** with respect to  $P$  and  $Q$  if  $\varphi(P) \subseteq Q$  (equivalently  $P = \varphi^{-1}(Q)$ ).

By the Theorem of the Primitive Element  $L = K(\alpha)$ .

Consider  $f = \text{MinPol}(\alpha | K)$ . Since  $\alpha \in R$ ,  $\varphi(f)$  has at least one root  $\beta$  in  $S$ ,

$$L := K(\alpha) \xleftrightarrow{\psi} \varphi(K)(\beta),$$

so there is at least one extension of  $\varphi$  from  $K$  to  $L$ .

Let  $\psi_1, \dots, \psi_r$  all such extensions of  $\varphi$  to  $L = K(\alpha)$ , and for a contradiction assume that none of them is order preserving with respect to  $Q = L \cap \sum R^2$ . Then  $\exists b_1, \dots, b_r \in L$ ,  $b_i > 0$  (in  $R$ ) and  $\psi_i(b_i) < 0$  (in  $S$ )  $\forall i = 1, \dots, r$ .

Consider  $L' := L(\sqrt{b_1}, \dots, \sqrt{b_r}) \subset R$ . Since  $[L : K] < \infty$ , also  $[L', K] < \infty$ .

So let  $\tau$  be an extension of  $\varphi$  from  $K$  to  $L'$ . In particular  $\tau|_L$  is one of the  $\psi_i$ 's. Say  $\tau|_L = \psi_1$ .

Now compute for  $b_1 \in L$ ,

$$\psi_1(b_1) = \tau(b_1) = \tau((\sqrt{b_1})^2) = (\tau(\sqrt{b_1}))^2 \in \sum S^2,$$

in contradiction with the fact that  $\psi_1(b_1) < 0$ . □

**Theorem 2.2.** *Let  $(K, P)$  be an ordered field and  $(R, \sum R^2)$  be a real closure of  $(K, P)$ . Let  $(S, \sum S^2)$  be a real closed field and assume that*

$$\varphi: (K, P) \hookrightarrow (S, \sum S^2)$$

*is an order preserving embedding. Then  $\varphi$  has a uniquely determined extension*

$$\psi: (R, \sum R^2) \hookrightarrow (S, \sum S^2).$$

*Proof.* Consider

$$\mathcal{L} := \{(L, \psi) : K \subset L \subset R; \psi: L \hookrightarrow S, \psi|_K = \varphi\}.$$

Let  $(L, \psi)$  be a maximal element. Then by Proposition 2.1 we must have  $L = R$ .

Therefore we have an order preserving embedding  $\psi$  of  $R$  extending  $\varphi$

$$\psi: R \hookrightarrow S.$$

We want to prove that  $\psi$  is unique. We show that  $\psi(\alpha) \in S$  is uniquely determined for every  $\alpha \in R$ .

Let  $f = \text{PolMin}(\alpha | K)$  and let  $\alpha_1 < \dots, \alpha_r$  all the real roots of  $f$  in  $R$ . Let  $\beta_1 < \dots < \beta_r$  be all the real roots of  $f$  in  $S$ . Since  $\psi: R \hookrightarrow S$  is order preserving, we must have  $\psi(\alpha_i) = \beta_i$  for every  $i = 1, \dots, r$ . In particular  $\alpha = \alpha_j$  for some  $1 \leq j \leq r$  and  $\psi(\alpha) = \beta_j \in S$ .  $\square$

**Corollary 2.3.** *Let  $(K, P)$  be an ordered field,  $R_1, R_2$  two real closures of  $(K, P)$ . Then exists a unique*

$$\varphi: R_1 \longrightarrow R_2$$

*$K$ -isomorphism (i.e. with  $\varphi|_K = \text{id}$ ).*

**Corollary 2.4.** *Let  $R$  be a real closure of  $(K, P)$ . Then the only  $K$ -automorphism of  $R$  is the identity.*

**Corollary 2.5.** *Let  $R$  be a real closed field,  $K \subseteq R$  a subfield. Set  $P := K \cap \sum R^2$  the induced order. Then*

$$K^{ralg} = \{\alpha \in R : \alpha \text{ is algebraic over } K\}$$

*is relatively algebraic closed in  $R$  and is a real closure of  $(K, P)$ .*

*Proof.* It is enough to show that  $K^{ralg}$  is real closed.

$K^{ralg}$  is real because  $Q := K^{ralg} \cap \sum R^2$  is an induced ordering.

Let  $a \in Q$ ,  $a = b^2$ ,  $b \in R$ . So  $p(x) = x^2 - a \in K^{ralg}[x]$  has a root in  $R$ .

One can see that  $b$  is algebraic over  $K$  (so  $b \in K^{ralg}$ ).

Similarly one shows that every odd polynomial with coefficients in  $K^{ralg}$  has a root in  $K^{ralg}$ .  $\square$

**Corollary 2.6.** *Let  $(K, P)$  be an ordered field,  $S$  a real closed field and  $\varphi: (K, P) \hookrightarrow S$  an order preserving embedding. Let  $L | K$  an algebraic extension. Then there is a bijective correspondence*

$$\{\text{extensions } \psi: L \rightarrow S \text{ of } \varphi\} \longrightarrow \{\text{extensions } Q \text{ of } P \text{ to } L\}$$

$$\psi \longmapsto \psi^{-1}(\sum S^2)$$

*Proof.*

( $\Rightarrow$ ) Let  $\psi: L \hookrightarrow S$  an extension of  $\varphi$ . Then indeed  $Q := \psi^{-1}(\sum S^2)$  is an ordering on  $L$ . Furthermore  $\psi^{-1}(\sum S^2) \cap K = \varphi^{-1}(\sum S^2) = P$ . So the extension  $\psi$  induces the extension  $Q$ .

( $\Leftarrow$ ) Conversely assume that  $Q$  is an extension of  $P$  from  $K$  to  $L$  ( $Q \cap K = P$ ). Note that if  $R$  is a real closure of  $(L, Q)$  then  $R$  is a real closure of  $(K, P)$  as well.

Now apply Theorem 2.2 to extend  $\varphi$  to  $\sigma: R \rightarrow S$ . Set  $\psi := \sigma|_L$  which is order preserving with respect to  $Q$ . So the map is well-defined and surjective. To see that it is also injective, assume

$$\psi_1: L \longrightarrow S, \quad \psi_2: L \longrightarrow S, \quad \psi_1|_K = \psi_2|_K = \varphi$$

which induce the same order

$$Q = \psi_1^{-1}(\sum S^2) = \psi_2^{-1}(\sum S^2)$$

on  $L$ . Let  $R$  be the real closure of  $(L, Q)$ . Apply Theorem 2.2 to  $\psi_1$  and  $\psi_2$  to get uniquely determined extensions

$$\sigma_1: R \longrightarrow S, \quad \sigma_2: R \longrightarrow S,$$

of  $\psi_1$  and  $\psi_2$  respectively.

But now  $\sigma_1|_K = \sigma_2|_K = \varphi$ . By the uniqueness part of Theorem 2.2 we get  $\sigma_1 = \sigma_2$  and a fortiori  $\psi_1 = \psi_2$ . □

**Corollary 2.7.** *Let  $(K, P)$  be an ordered field,  $R$  a real closure,  $[L : K] < \infty$ . Let  $L = K(\alpha)$ ,  $f = \text{MinPol}(\alpha | K)$ . Then there is a bijection*

$$\{\text{roots of } f \text{ in } R\} \longrightarrow \{\text{extensions } Q \text{ of } P \text{ to } L\}.$$

*Proof.* If  $\beta$  is a root we consider the  $K$ -embedding

$$\varphi_\alpha: L \hookrightarrow R$$

such that  $\varphi_\alpha(\alpha) = \beta$ . Set  $Q := \varphi^{-1}(\sum R^2)$  ordering on  $L$  extending  $P$ . □

**Example 2.8.**  $K = \mathbb{Q}(\sqrt{2})$  has 2 orderings  $P_1 \neq P_2$ , with  $\sqrt{2} \in P_1$ ,  $\sqrt{2} \notin P_2$ . The Minimum Polynomial of  $\sqrt{2}$  over  $\mathbb{Q}$  is  $p(x) = x^2 - 2$ .